IN THE DRAWINGS:

A duplicate drawing is submitted herewith. The color values set forth in the disclosure are accurate. The color photograph is produced as best as possible to reflect those values.

RESPONSE UNDER 37 CFR 1.105

The undersigned is advised that the subject variety was first offered for sale in Germany in 1998. As of the United States filing date, no sales were made nor was product available in the United States. This variety was the subject of Furopean Union Plant Breeder's Rights Certificate No. 7038. A copy of this Certificate is currently being located and will be forwarded to the United States Patent and Trademark Office upon location.

REMARKS

The Office Action of December 4, 2001 has been reviewed and the Examiner's comments carefully considered. The present Amendment provides a substitute specification addressing the outstanding issues in the present application. Applicant notes that two copies of the drawing are required, and, accordingly, please find a duplicate drawing attached hereto.

The Examiner has objected to the disclosure, pursuant to 37 C.F.R. § 1.163(a) and under 35 U.S.C. § 112, first and second paragraphs. Applicant believes that the amendments made to the specification specifically address the Examiner's objections detailed in Paragraphs A-K of the Office Action. Withdrawal of the objections and reconsideration of the rejection under 35 U.S.C. § 112, first and second paragraphs, is respectfully requested.

Reconsideration of the rejection, withdrawal of the objections, and allowance of the claim are respectfully requested.

Respectfully submitted,

WEBB ZIESENHEIM LOGSDON ORKIN & HANSON, P.C.

Ву

Russell D. Orkin

Registration No. 25,363

Attorney for Applicant

436 Seventh Avenue

700 Koppers Building

Pittsburgh, PA 15219-1818 Telephone: 412-471-8815

Facsimile: 412-471-4094

E-mail: webblaw@webblaw.com



GERANIUM PLANT NAMED 'TIKVIO'

CLASSIFICATION

Pelargonium x hortorum (zonal geranium)

VARIETAL DENOMINATION

'Tikvio'

BACKGROUND OF THE INVENTION

The present invention comprises a new and distinct cultivar of *Pelargonium x hortorum* known by the commercial names 'pac[®] Antik Violet, Tikvio^(s) and pac[®] Pillar Violet, Tikvio^(s) (Breeder No. P-6138). The new variety was discovered in 1996 in a selected breeding program in Dresden, Germany. The new variety is a selection a population of seedlings resulting from breeding and crossing of diploid group 95K-37. The new variety is vigorous with long internodes similar to the plants of its heritage but differs in its bright violet flower color. The new variety was first asexually reproduced by cuttings in Dresden, Germany. The new variety has been trial and field tested at Connellsville and has been found to retain its distinctive characteristics and remain true to type through successive propagations.

The new variety is comparable to 'Tikorg' (U.S. Patent Application Serial No. 09/753,976) but has less branching, violet flower color instead of orange and is hairier.

The cultivar, when grown in a glass greenhouse in Connellsville, Pennsylvania using natural light and temperatures of 60°F night and 68°F day, has a response time of six weeks from a rooted cutting to flowering in a 2 inch pot and a response time of 10-12 weeks to a saleable pillar product. The response time was determined on plants grown in soilless media employing constant fertilizer 200-250 parts per million of nitrogen and potassium in full light.

DESCRIPTION OF THE DRAWINGS

The accompanying photographic drawing illustrates the new variety, with the color being as nearly true as is possible with color illustrations of this type.

DESCRIPTION OF THE PLANT

The following detailed description sets forth the characteristics of the

readings were taken indoors under 200-220 foot candles of fluorescent cool white

Market 15: and

light. Color references are primarily to the R.H.S. Colour Chart of The Royal Horticultural Society of London, 1995 edition.

PLANT

[Classification:

Botanical:

Pelargonium x hortorum.

Commercial: Zonal geranium.]

Form:

Tall bush.

Height:

24-36 inches.

Growth:

Free branching, lateral to upright growth.

Strength:

If grown in pillar form, support required.

Foliage:

Stalked leaf.

Rooting time:

14-21 days depending on conditions.

Leaves:

Size:

Length:

5.5-7.0 cm.

Width:

7.5-8.0 cm.

Shape:

Reniform; cordate base.

Margin:

Crenate.

Texture:

Pubescent.

Color:

Upper surface:

Green Group 137A.

Lower surface:

Green Group 137C.

Zone:

Greyed-Purple Group 187A.

Ribs and veins:

Venation:

Palmate.

Color:

Yellow-Green Group 144D.

Petioles:

Color:

Yellow-Green of Group 144A with shading some

Greyed-Purple Group 187B.

Stem:

Color:

Green Group 137C.

Internode Length:

2.5-4.5 cm.

Diameter:

8 mm.

THE BUD

Shape when just beginning to show color:

Overall:

Semi-spherical.

Individual bud:

Elongated elliptical (oval).

Size when just beginning to show color:

Overall:

4.0-4.5 cm across.

Individual bud:

0.5-0.7 cm wide; 1.2-1.5 cm long.

Number of buds per cluster: 15-20.

INFLORESCENCE

Blooming habit:

Continuous flowering.

Size:

11.5-12.5 cm across.

Borne:

Flower on umbel; umbel on pedicel; pedicel on peduncle.

Flowers:

Form:

Slightly cupped, some petals overlap.

Number of flowers per umbel: 20-25.

Size:

Diameter:

4.5-4.7 cm.

Depth:

1-1.5 cm.

Petals:

Color:

Upper surface:

Red-Purple Group 66A, some petals have slight outer edge rim of Red-Purple Group 57A. Toward the base of petal the color shades to Red Group 50A. On the upper two petals there is an area of

Red Group 52A shading to small areas of

Lower surface:

Wide color range including: Red-Purple

Group 64C, Red-Purple Group 63B,

Red-Purple Group 61C, Red-Purple

Groups 58B and 58C, and Red Group

50B. Some petals have dark veining that

shows through.

Shape:

Round at top with a narrower bottom.

Size:

Length:

2.4-2.6 cm.

Width:

1.8-2.4 cm.

Margin:

[Complete] Bicrenate.

Number per flower:

5.

Sepals:

Color:

Green Group 138A.

Size:

8 mm long; 5 mm wide.

Number per flower: 5.

Texture and appearance:

Smooth, highly reflective.

Tonality from a distance:

Striking magenta flowers with tones of deep red, and

violet varying in amount and intensity.

Petaloids:

None.

Pedicel:

Length:

4.0-4.3 cm.

Diameter:

1.0-1.5 mm.

Color:

Greyed-Purple Group 184A with Yellow-Green Group 144D at

base.

Peduncle:

Length:

18-20 cm.

Diameter:

4-5 mm.

Color:

Yellow-Green Group 144A with shading of Greyed-Purple

Group 187B.

REPRODUCTIVE ORGANS

Stamens:

Anthers:

0.2-0.25 cm.

Filaments:

Length:

0.4-0.5 cm.

Color:

[White with magenta tips] Yellow-Green Group 145C.

Pollen:

[Orange] Orange Group 24B.

Pistils:

Number:

One.

Length:

0.7 cm.

Stigma:

Number:

One.

Color:

[Red-Violet] Yellow-Green Group 150D.

Style:

Length:

0.2-0.3 cm.

Color:

[Red-Violet] Yellow-Green Group 150D.

Ovaries:

Length:

0.5 cm.

Width:

0.3 cm.

Color:

[Green] Green Group 142B.

Fruit: None observed.

GENERAL CHARACTERISTICS

- 1. 'Tikvio' is a very vigorous grower which allows it to be trained as a pillar.
- 2. 'Tikvio' is very floriferous and unique to the geranium industry.
- 3. Tolerant to drought and some heat; not winter hardy.